

# Regional ventilation/perfusion mismatch pattern in patient with Swyer James (MacLeod's) syndrome

Sait Sager, Sertac Asa, Reşit Akyel, Ersan Atahan<sup>1</sup>, Bedii Kanmaz

Departments of Nuclear Medicine and <sup>1</sup>Chest Disease, Cerrahpaşa Medical Faculty, Istanbul University, Cerrahpasa, Fatih, Istanbul, Turkey

Swyer James (McLeod's) syndrome (SJMS) is an uncommon disease, which occurs as a result of childhood bronchiolitis obliterans. Patients may not be diagnosed until later in their life. A 46-year-old man underwent ventilation/perfusion scintigraphy for acute onset of dyspnea. The scan showed markedly diminished ventilation and perfusion unilaterally on the right middle and inferior lobes. However, mismatched ventilation-perfusion pattern was shown on the upper right lobe, which was consistent with pulmonary embolism. Unilaterally matched ventilation/perfusion defect can be seen in SJMS in lung scintigraphy; however, when pulmonary embolism may accompany, scintigraphy should be carefully examined.

**Key words:** McLeod's syndrome, pulmonary embolism, scintigraphy, Swyer James

**How to cite this article:** Sager S, Asa S, Akyel R, Atahan E, Kanmaz B. Regional ventilation/perfusion mismatch pattern in patient with Swyer James (MacLeod's) syndrome. *J Res Med Sci* 2014;19:904-6.

## INTRODUCTION

Swyer James (McLeod's) syndrome (SJMS) is a rare condition (was seen 0.01% in 17459 cases) characterized by unilateral hyperlucency of a part of or the entire lung, which was first described by Swyer and James than MacLeod.<sup>[1,2]</sup> First, this syndrome was thought to be congenital in origin and was attributed to hypoplasia of the pulmonary artery. SJMS is a variant of postinfectious bronchiolitis obliterans that is most commonly the sequela of a viral (adenovirus or RVS) infection during infancy or early childhood. Mycoplasma infection has also been implicated.<sup>[3]</sup> SJMS may be seen following measles, pertussis, tuberculosis, and mycoplasma infections.<sup>[4]</sup>

## CASE REPORT

A 46-year-old male patient was admitted to Istanbul University, Cerrahpasa Medical Faculty, Department of Chest Disease, Istanbul in 2009 with dyspnea and chest pain. While examination of the patient it is learned that he had a history of multiple respiratory infections in childhood. Chest examination showed diminished breath sounds over the right hemithorax. Chest X-ray (CXR) showed hyperlucency on the lower part of the right hemithorax [Figure 1].

Thorax computed tomography (CT) revealed bronchiectatic changes, hyperlucency of the lower

part of the right hemithorax due to hypoplastic left pulmonary artery [Figure 2]. Thus, diagnosis of SJMS associated with right lung middle lobe hypoplasia was made.


Patient referred to the nuclear medicine department for ventilation-perfusion scintigraphy to rule out pulmonary thromboembolism. Ventilation/perfusion scintigraphy (VQ) was performed in the same day. Technetium-99m-labeled macroaggregated human albumin (99mTc-MAA) perfusion scan was performed after the injection of 185-200 MBq of 99mTc-MAA into an antero-cubital vein, and Tc-99m labeled carbon particles (Technegas) was performed after inhalation of technegas with tidal breathing in upright position for ventilation scintigraphy. Planar images and single photon emission computed tomography images of each scan were obtained by a double-headed gamma camera. VQ images showed severe unilaterally matched VQ defects on the right middle and inferior lobes. However, mismatched ventilation-perfusion pattern was shown on the upper right lobe, which was consistent with pulmonary embolism [Figure 3].

## DISCUSSION


Swyer-James-MacLeod syndrome is a rare lung disorder, which occurs as a result of childhood bronchiolitis obliterans. The onset of symptoms typically occurs during infancy or early childhood

**Address for correspondence:** Dr. Sait Sager, Department of Nuclear Medicine, Cerrahpasa Medical Faculty, Istanbul University, Cerrahpasa, Fatih, Istanbul, Turkey. E-mail: saitsager@yahoo.com


**Received:** 02-11-2013; **Revised:** 22-05-2014; **Accepted:** 27-05-2014



**Figure 1:** Chest X-ray showed hyperlucency on the lower part of the right hemithorax



**Figure 2:** Computed tomography revealed bronchiectatic changes, hyperlucency of lower part of the right hemithorax



**Figure 3:** Posterior, right posterior oblique (RPO) and right lateral perfusion images (top row) and posterior, RPO, right lateral ventilation images (lower row) showed unilaterally matched VQ defects of the lower part of the right lung

in association with frequent respiratory infections. Patients usually present with productive cough, shortness of breath, and dyspnea on exertion. In SJMS,

involved lung or portion of the lung does not grow normally and is slightly smaller than the opposite lung. The characteristic radiographic appearance is that of pulmonary hyperlucency, caused by over distention of the alveoli in conjunction with diminished arterial flow. In general, this disorder is diagnosed in childhood after evaluations for recurrent respiratory infections.<sup>[5]</sup> After the initial infective insult, radiographic findings appear after months to years. Usual investigations performed are chest radiograph, high-resolution CT, magnetic resonance imaging, angiography, ventilation-perfusion scanning.

Swyer-James-MacLeod syndrome is usually diagnosed with chest radiography which shows hyperlucency of a lung lobe and a small hemithorax on the affected side. However, CXR can be insensitive and may appear normal. Despite characteristic findings by chest radiography, CT is the most valuable imaging technique in the diagnosis of SJMS. When using CT scan peribronchial thickening and the diagnosis of bullae and subsequent air trapping can be seen.<sup>[3]</sup>

Swyer-James-MacLeod syndrome is often referred to as hyper- or trans-lucent unilateral lung because of the decreased parenchymal perfusion.<sup>[6]</sup> Although one lung or one lobe can be affected, CT has made it apparent that the disease may be more heterogeneous in distribution and contralateral parenchymal lesion may be present.<sup>[7]</sup>

Ventilation/perfusion scintigraphy is a diagnostic modality that performed to rule out pulmonary embolism and shows a decrease in ventilation to the affected lung, secondary to emphysematous changes and a matched decrease in perfusion.<sup>[8]</sup> In our patient, VQ scintigraphy showed matched defect in the affected right middle and lower lobe.

Pulmonary angiography is extremely valuable in showing narrowed and withered tree-like pulmonary arteries in the affected side. Magnetic resonance (MR) angiography demonstrates smaller pulmonary artery and its branches on the affected side. Serdengecti *et al.* compared MR angiography with ventilation-perfusion scintigraphy in 11 patients and concluded that the magnetic resonance angiography is a fast, accurate, without radiation, and noninvasive technique for supporting the diagnosis of Stevens-Johnson syndrome.<sup>[9]</sup>

The important differential diagnosis of the SJMS are congenital lobar emphysema, congenital hypoplasia of the lung, compensatory unilateral emphysema secondary to lobectomy, pulmonary embolic disease, pneumothorax, and foreign body in air way.

## AUTHORS' CONTRIBUTIONS

All authors have contributed in designing and conducting the study. All authors have assisted in preparation of the first draft of the manuscript or revising it critically for important intellectual content. All authors have read and approved the content of the manuscript and confirmed the accuracy or integrity of any part of the work.

## REFERENCES

1. Swyer PR, James GC. A case of unilateral pulmonary emphysema. *Thorax* 1953;8:133-6.
2. Macleod WM. Abnormal transradiancy of one lung. *Thorax* 1954;9:147-53.
3. Lucaya J, Gartner S, García-Peña P, Cobos N, Roca I, Liñan S. Spectrum of manifestations of Swyer-James-MacLeod syndrome. *J Comput Assist Tomogr* 1998;22:592-7.
4. Dillman JR, Sanchez R, Ladino-Torres MF, Yarram SG, Strouse PJ, Lucaya J. Expanding upon the unilateral hyperlucent hemithorax in children. *Radiographics* 2011;31:723-41.
5. Fishman AP. *Pulmonary Disease and Disorders*. 2<sup>nd</sup> ed. New York: McGraw-Hill; 1988. p. 494, 2550-1.
6. Salmanzadeh A, Pomeranz SJ, Ramsingh PS. Ventilation-perfusion scintigraphic correlation with multimodality imaging in a proven case of Swyer-James (Macleod's) syndrome. *Clin Nucl Med* 1997;22:115-8.
7. Ghossain MA, Achkar A, Buy JN, Rochemaure J, Vadrot D. Swyer-James syndrome documented by spiral CT angiography and high resolution inspiratory and expiratory CT: An accurate single modality exploration. *J Comput Assist Tomogr* 1997;21:616-8.
8. Bajc M, Neilly JB, Miniati M, Schuemichen C, Meignan M, Jonson B, *et al.* EANM guidelines for ventilation/perfusion scintigraphy: Part 1. Pulmonary imaging with ventilation/perfusion single photon emission tomography. *Eur J Nucl Med Mol Imaging* 2009;36:1356-70.
9. Serdengecti M, Sakarya ME, Ilerisoy ZY, Odev K. Comparison of ventilation-perfusion scintigraphy with MR angiography in patients with Swyer-James syndrome. *Clin Nucl Med* 2013;38:237-40.

**Source of Support:** Nil, **Conflict of Interest:** None declared.