

# Antibiotic resistance: A disregarded concern for misuse of azithromycin in COVID-19 treatment

Dear Editor,


In late November 2019, a new viral infection began to spread from Wuhan, China, resulting in an outbreak that was later declared a pandemic on March 12, 2020. So far, over 125,000,000 confirmed cases and 2,700,000 deaths have resulted from the spread of coronavirus disease-2019 (COVID-19) infection. Concerns about the disease increase when we consider the fact that no definitive cure is currently available. Due to time constraints, many researchers and physicians have begun to use existing drugs that could rationally invade the underlying mechanisms of the disease, hoping to achieve positive results in stopping the progression of the disease or preventing the expected complications.<sup>[1]</sup> One of the proposed protocols was the use of azithromycin, mostly combined with hydroxychloroquine. At the beginning of the pandemic, azithromycin was thought to be a beneficial agent against COVID-19. However, at present, it is not recommended – either alone or in combination with hydroxychloroquine – since it has shown no antiviral effect, and no positive results have been observed on patients' mortality or the severity of the disease.<sup>[2]</sup>

The recommenders used to emphasize the fact that the usefulness and efficiency of macrolides have been previously proven in similar viral cases such as

rhinovirus infection and influenza.<sup>[3]</sup> Furthermore, in addition to the antibacterial properties, macrolides have shown immunomodulatory effects, including anti-inflammatory outcomes. Macrolides' ability to suppress the infections lies in their ability to terminate the viral replication, block the internalization of the virus into the host cells, and modify the endosomal and lysosomal pH.<sup>[4]</sup>

A wide range of usage and fewer side effects, along with the possibility of rapid production, has turned azithromycin into a strategic antimicrobial agent. Despite all potential benefits, it should be noted that considering azithromycin as a treatment for COVID-19 infection can have its consequences. According to the World Health Organization, azithromycin is one of the "highest priority critically important antimicrobials," which should be used prudently to prevent antimicrobial resistance.<sup>[5]</sup> Therefore, the excessive consumption of azithromycin – among the terrified community, who are looking for ways to prevent the infection, and the clinical wards treating the COVID-19 patients – may result in an undeniable crisis of resistance to this irreplaceable agent. This could also leave short-term complications if the resistance happens in vulnerable COVID-19 patients with immunodeficiency or underlying diseases, such as cancer and metabolic disorders. Hence, misuse and overuse of azithromycin have to be strongly avoided, and it should always be administered with caution in vulnerable patients [Figure 1].

As a global state of emergency, any effort to achieve a reliable treatment or prevention is quite rational and valuable during the pandemics. However, it should be considered that a wrong move may result in an irreversible catastrophe, which can cause life-long trouble for human beings.



**Figure 1:** Status of viral (SARS-CoV-2) and bacterial (e.g., *Streptococcus pneumoniae*) resistance. The blue color indicates susceptibility and the red color indicates resistance. At the beginning, azithromycin is prescribed in combination with some antiviral agents. Azithromycin-resistant bacteria appear and gradually grow in number. Eventually, azithromycin loses its place as an effective antibiotic. By the spread of the resistant bacteria, the infected patients with bacterial pneumonia are now more susceptible to other infections, including viral pneumonia, which results in fatal superinfections

**Financial support and sponsorship**

Nil.

**Conflicts of interest**

There are no conflicts of interest.

**Farid Jahanshahlo<sup>1</sup>, Mohammad-Salar Hosseini<sup>1,2,3</sup>**

<sup>1</sup>Student Research Committee, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran, <sup>2</sup>Research Center for Evidence-Based Medicine, Tabriz University of Medical Sciences, Tabriz, Iran, <sup>3</sup>Emergency Medicine Research Team, Tabriz University of Medical Sciences, Tabriz, Iran

**Address for correspondence:** Dr. Mohammad-Salar Hosseini, Research Center for Evidence-Based Medicine, Tabriz University of Medical Sciences, Golgasht Street, Postal Code 5166/15731, Tabriz, EA, Iran.  
E-mail: hosseini.msalar@gmail.com

**Submitted:** 20-Sep-2020; **Revised:** 28-Mar-2021;

**Accepted:** 14-Apr-2021; **Published:** 18-Oct-2021

**REFERENCES**

1. Madrid PB, Panchal RG, Warren TK, Shurtleff AC, Endsley AN, Green CE, *et al.* Evaluation of ebola virus inhibitors for drug repurposing. *ACS Infect Dis* 2015;1:317-26.
2. Siemieniuk RA, Bartoszko JJ, Ge L, Zeraatkar D, Izcovich A, Kum E, *et al.* Drug treatments for COVID-19: Living systematic review and network meta-analysis. *BMJ* 2020;370:m2980.

3. Lee N, Wong CK, Chan MCW, Yeung ESL, Tam WWS, Tsang OTY, *et al.* Anti-inflammatory effects of adjunctive macrolide treatment in adults hospitalized with influenza: A randomized controlled trial. *Antiviral Res* 2017;144:48-56.
4. Ohe M, Shida H, Jodo S, Kusunoki Y, Seki M, Furuya K, *et al.* Macrolide treatment for COVID-19: Will this be the way forward? *Biosci Trends* 2020;14:159-60.
5. World Health Organization. Highest Priority Critically Important Antimicrobials; 2019. Available from: <https://www.who.int/foodsafety/cia/en/>. [Last accessed on 2020 Sep 12].

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**Access this article online**

Quick Response Code:



Website:

[www.jmsjournal.net](http://www.jmsjournal.net)

DOI:

10.4103/jrms.JRMS\_1124\_20

**How to cite this article:** Jahanshahlo F, Hosseini MS. Antibiotic resistance: A disregarded concern for misuse of azithromycin in COVID-19 treatment. *J Res Med Sci* 2021;26:101.

© 2021 Journal of Research in Medical Sciences | Published by Wolters Kluwer - Medknow