

Estimating the best fraction of inspired oxygen for calculation of PaO₂/FiO₂ ratio in acute respiratory distress syndrome due to COVID-19 pneumonia

Leila Kadkhodai^{1,2}, Mahmoud Saghaei^{1,2}, Mohammadreza Habibzadeh^{1,2}, Babak Alikiaii^{1,2}, Seyed Jalal Hashemi^{1,2}

¹Department of Anesthesia, Isfahan University of Medical Sciences, Isfahan, Iran, ²Anesthesiology and Critical Care Research Center, Isfahan University of Medical Sciences, Isfahan, Iran

Background: The ratio of partial pressure of oxygen in arterial blood (PaO₂) to the fraction of inspiratory oxygen concentration (FiO₂) is an indicator of pulmonary shunt fraction. PaO₂/FiO₂ (P/F) ratio is used to classify severity of acute respiratory distress syndrome (ARDS). With the same shunt fraction, P/F ratio decreases with increases in FiO₂ which may lead to errors in classifying severity of ARDS. The effect of FiO₂ on P/F ratio has not been investigated in COVID-19 pneumonia. In this study, we estimated the best FiO₂ for the calculation of P/F ratio in a sample of patients with ARDS due to COVID-19 pneumonia. **Materials and Methods:** Blood gas and ventilatory data of 108 COVID-19 ARDS patients were analyzed in a cross-sectional observational study. Using Oxygen Status Algorithm the calculated shunt fraction served a basis for calculating P/F ratio for different FiO₂. The severity of ARDS determined by P/F ratios at each FiO₂s was compared with the shunt-based severity to find the optimum FiO₂ for calculation of P/F ratio so the resulting classification has the best match with the reference classification. **Results:** A FiO₂ of 1.0 for calculation of P/F ratio and ARDS classification showed the best match with shunt-based ARDS classification. A regression model was obtained with the PaO₂, patient's original FiO₂, Hemoglobin concentration, and SaO₂ as the independent predictors of the P/F ratio for the FiO₂ of 1.0. **Conclusion:** This study shows a FiO₂ of 1.0 as the best value for correct calculation of P/F ratio and proper classification of ARDS.

Key words: Acute respiratory distress syndrome, COVID-19, mechanical ventilation, oxygenation indices, P/F ratio, pulmonary shunt

How to cite this article: Kadkhodai L, Saghaei M, Habibzadeh M, Alikiaii B, Hashemi SJ. Estimating the best fraction of inspired oxygen for calculation of PaO₂/FiO₂ ratio in acute respiratory distress syndrome due to COVID-19 pneumonia. *J Res Med Sci* 2022;27:38.

INTRODUCTION

A fundamental step in the management of patients with arterial hypoxia is to assess the extent of damage to the gas exchange function of the lungs. The measurement of pulmonary shunt fraction which is defined as the fraction of mixed venous blood bypassing oxygenation in the lung capillaries is the most accurate method to quantify the extent of pulmonary damage.^[1]

The reason why shunt fraction is the best indicator of lung involvement associated with oxygenation

disturbances lays in its independence from inspired oxygen concentration and the type of oxygenation deficit.^[2] Clinical measurement of shunt fraction requires insertion of pulmonary artery catheter to obtain mixed venous blood sample from pulmonary artery, which unfortunately is a risky procedure especially in critically ill patients and therefore is not feasible under most clinical conditions.^[3] Therefore, clinicians use the ratio of oxygen partial pressure in arterial blood (PaO₂) to inspiratory fraction of oxygen (FiO₂) as a surrogate for pulmonary shunt fraction.^[4] In healthy controls, PaO₂/FiO₂ (P/F) ratio varies from 400 to 500 at sea level and reduces with decreasing barometric pressure.^[5,6] The

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

Access this article online

Quick Response Code:



Website:

www.jmsjournal.net

DOI:

10.4103/jrms.jrms_558_21

Address for correspondence: Dr. Mahmoud Saghaei, Al-Zahra Medical Center, Isfahan, Iran.

E-mail: mahmood.saghaei@gmail.com

Submitted: 30-Jun-2021; **Revised:** 24-Nov-2021; **Accepted:** 14-Dec-2021; **Published:** 30-May-2022

Berlin definition uses P/F ratio for classification of acute respiratory distress syndrome (ARDS) into mild ($200 < P/F \text{ ratio} \leq 300$), moderate ($100 < P/F \text{ ratio} \leq 200$), and severe ($P/F \text{ ratio} \leq 100$) form of the syndrome.^[7] Unfortunately with the same value of shunt fraction the P/F ratio varies with the value of FiO₂ at which the PaO₂ is measured.^[8] Therefore its value as a surrogate for pulmonary shunt fraction and as a discriminating factor for ARDS classification significantly deteriorates with changes in the FiO₂. For example, at a fixed pulmonary shunt of 40% the P/F ratio would be 220 (mild ARDS) and 64 (severe ARDS) at FiO₂ of 0.21 and 1.0, respectively.^[9]

A previous study on ARDS, showed considerable discrepancies in severity categorization using P/F ratios corrected for FiO₂ compared to noncorrected values.^[10] The effect of FiO₂ on P/F ratio has not been investigated previously. Since FiO₂- PaO₂ relationship may be different in COVID-19 induced pneumonia it's necessary to investigate the effects of FiO₂ on P/F ratio and the consequences on ARDS classification in ARDS cases due to COVID-19 pneumonia. COVID-19 induced pneumonia and ARDS has been a major challenge since the start of the pandemics and necessitates special attention.^[11,12] In this study on a sample of adult patients with COVID-19-induced ARDS patients, we used blood gas and ventilatory data of patients together with Oxygen Status Algorithm (OSA) to calculate P/F ratios for a range of FiO₂s and compared the resultant severity categorization with the severity classification defined by the estimated shunt fraction to obtain the optimum FiO₂ for calculation of P/F ratio.

MATERIALS AND METHODS

Ethics committee of Research Department, Isfahan University of Medical Sciences approved the study (IR. MUI. MED. REC.1399.1010). In this cross-sectional observation study, blood gas and ventilatory data collected from mechanically ventilated patients with COVID-19-induced ARDS. Pregnant patients, patients with cardiovascular disorders, history of chronic respiratory disease, heavy smoking, history of mechanical ventilation in the last year, and those on drugs influencing pulmonary shunt (hydralazine, dopamine, dobutamine, and nitrate-containing substances) excluded from the study.

All patients were under mechanical ventilation using SIMV mode with lung protecting strategy. A tidal volume = 4–6 ml/kg, positive end-expiratory pressure = 5–10 cmH₂O, and respiratory rate manipulated according to the level of arterial partial pressure of CO₂.

Oxygen status algorithm

OSA version 3.0 is a Windows program developed by Siggaard-Andersen and coworkers.^[13] OSA estimates the

acid-base and oxygen status of the blood and displays the results in the form of charts, graph, blood gas map, and diagram. It also allows changing some respiratory parameters and studying the effects on others parameter while keeping some parameters constant. This application has been used extensively as a laboratory, research, and clinical tools for studying and managing acid-base and oxygen status of patients, especially in the intensive care unit. Blood gas and ventilatory data of the patient must be entered directly into the program using main program window or inputted as a data file. The barometric pressure of the area must be entered into the program and the program corrects all calculated values for the barometric pressure at the sea level (760 mmHg).^[14] We entered patient blood gas and ventilatory data into the application, and then recorded the shunt fraction calculated by the program. In the next step, we kept shunt fraction constant in the program and changed FiO₂ from 0.2 to 1.0 in increments of 0.1. After each change in FiO₂, the calculated values for PaO₂ and the resulting P/F ratio were recorded. Therefore for each patient, nine different values of P/F ratio for different values of FiO₂ were simulated.

Severity categorization

For each estimated P/F ratio obtained from OSA the severity of ARDS was determined based on the threshold values recommended by Berlin definition task force^[7] as shown in Table 1. As the reference classification, the severity of ARDS in each patient was determined by using the threshold values for shunt fraction defined in the Berlin definition specifications [Table 1].^[7]

Statistical analysis

A sample size of 84 was calculated using the following formula for a confidence level of 80%, a margin of error equal to 5% using previously reported value of 70 as the P/F ratio standard deviation (s) (SD)^[2,15] to estimate the mean P/F ratio with an error of 15.

$$N = \frac{\left(Z_{1-\frac{\alpha}{2}} \right)^2 \times S^2}{d^2} = \frac{(1.96)^2 \times 70^2}{15^2} = 83.66 \approx 84$$

Table 1: Thresholds of PaO₂/FiO₂ ratio and shunt fractions for classification of acute respiratory distress syndrome

ARDS category	Definition
No ARDS	P/F ratio > 300 Shunt < 16%
Mild	200 < P/F ratio ≤ 300 16% ≤ shunt < 26
Moderate	100 < P/F ratio ≤ 200 26% ≤ shunt < 32
Severe	P/F ratio ≤ 100 Shunt ≥ 32%

P/F ratio = PaO₂/FiO₂ ratio; ARDS = Acute respiratory distress syndrome

Resulting classifications for each value of P/F ratio associated with each hypothetical FiO₂ were compared with reference classification based on shunt fraction to calculate the proportions of correct and wrong classifications. The proportions of correct and wrong classifications for each FiO₂ were compared with each other using Chi-square statistics together with Spearman correlations to measure the association between P/F ratio-based classifications and reference classification. Multiple linear regression analysis used to build a model for predicting simulated P/F ratio based on blood gas data. We used SPSS version 26 for statistical analysis. A $P < 0.05$ was considered statistically significant. Data were presented as mean \pm SD or n (%) where applicable. The SPSS data were exported as comma-delimited file which used by a python script to produce appropriate data files for input into OSA application.

RESULTS

A total of 108 (51 male and 48 female) 20–90 years' old (61.8 ± 13.7) cases were studied. Data from nine cases had missing values and therefore excluded from the analysis. PaO₂ ranged from 30 to 90 mmHg (66 ± 24), FiO₂ from 0.30 to 1.0 (0.76 ± 0.24), P/F ratio from 30 to 428 (109 ± 86). Demographic data and clinically measured ventilatory parameters are summarized in Tables 2 and 3 which shows a wide range of shunt fractions and P/F ratios.

Calculated pulmonary shunt for each patient ranged from 1.35% to 69.5% (34 ± 15.6).

Using shunt-based severity as the reference for severity classification showed that the best match belonged to P/F ratio for an FiO₂ of 1.0 (P/F_{1.0}) [Figure 1, and Table 4]. Using an FiO₂ of 1.0, severity of the ARDS classified correctly for 84 patients (84.8%, 95% confidence interval 76–91, Spearman correlation[®] of 0.944, $P < 0.001$). The worst classification was with FiO₂ of 0.20 with only 26 patients classified




Figure 1: Proportions of correct and wrong ARDS classifications among different types of P/F ratios. ARDS = Acute Respiratory Distress Syndrome; FiO₂ = Fraction of Inspiratory Oxygen

correctly (26.3, 95% confidence interval [CI], 18–36, $r = 0.722$, $P < 0.001$).

Multiple regression analysis with P/F_{1.0} as dependent variable showed PaO₂, patient's original FiO₂, hemoglobin concentration (Hb), and oxygen saturation of hemoglobin (SaO₂) as the independent predictors ($R^2 = 0.882$, $P = 0.000$). Regression coefficients for these predictor variables yielded the following model:

$$P/F_{1.0} = 393 + 5.426 \times PaO_2 - 237.2 \times FiO_2 - 8.134 \times Hb - 3.769 \times SaO_2$$

Table 2: Different demographic, blood gas, and ventilatory data of the patients (n=99)

Variable	Value
Male	51 (51.5)
Female	48 (48.5)
Age (years)	61.8 \pm 13.7 (20-90)
PaO ₂ (mmHg)	66 \pm 24 (30-90)
SaO ₂ (%)	87 \pm 9.6 (51-97)
FiO ₂	0.76 \pm 0.24 (0.30-1.0)
PaO ₂ /FiO ₂	109 \pm 86 (30-428)
PaCO ₂ (mmHg)	53 \pm 18.5 (19-135)
pH	7.30 \pm 0.08 (7.06-7.59)
Hemoglobin (g/dL)	11 \pm 2.3 (5.1-18.2)
Pulmonary shunt (%)	34 \pm 15.6 (1.35-69.5)
Static compliance (ml/cmH ₂ O)	41 \pm 29 (5-150)
Airway resistance (cmH ₂ O/L/s)	10.4 \pm 6 (1-34)
Peak inspiratory pressure (cmH ₂ O)	28 \pm 8.4 (11-50)
Positive end-expiratory pressure (cmH ₂ O)	8.8 \pm 3.4 (5-25)
Mean airway pressure (cmH ₂ O)	15.7 \pm 4.7 (6-27)
Minute ventilation (L/min)	11 \pm 2.9 (4.4-17.5)
Total respiratory rate (breath/min)	25 \pm 7.7 (10-54)
Mechanical ventilation mode	
SIMV	81 (81.8)
A/C	10 (10.1)
SPV	8 (8.1)

Data are mean \pm SD (minimum-maximum) or n (%) where applicable. SD=Standard deviation; SIMV=Synchronized Intermittent Mandatory Ventilation; A/C=Assist/Control; PSV=Pressure Support Ventilation

Table 3: Estimated PaO₂/FiO₂ ratio and proportions of correct classifications associated with each value of PaO₂/FiO₂ ratio

FiO ₂	Mean \pm SD (range) ^a	Correct classifications (%)	Spearman correlation ^b
0.2	235 \pm 74 (50-480)	26 (26.3, 18-36)	0.722
0.3	193 \pm 83 (88-528)	38 (38.4, 29-49)	0.823
0.4	165 \pm 97 (69-564)	39 (39.4, 30-50)	0.677
0.5	150 \pm 112 (56-591)	56 (56.6, 46-67)	0.732
0.6	144 \pm 126 (47-611)	73 (73.7, 64-82)	0.862
0.7	141 \pm 138 (41-626)	79 (79.8, 71-87)	0.934
0.8	142 \pm 148 (37-638)	76 (76.8, 67-85)	0.903
0.9	146 \pm 157 (34-656)	82 (82.8, 74-90)	0.912
1.0	152 \pm 165 (31-653)	84 (84.8, 76-91)	0.944

Data are mean \pm SD or n (%) 95% CI. ^aAll P/F ratios calculated using oxygen status algorithm corrected to a barometric pressure of 760 mmHg (11). ^bAll correlations were significant at $P < 0.001$ value. P/F ratio: PaO₂/FiO₂ ratio. SD=Standard deviation; CI=Confidence interval

Table 4: Proportions of acute respiratory distress syndrome severity based on shunt fraction and PaO₂/FiO₂ ratio with FiO₂ of 1.0

Shunt fraction	P/F _{1.0}				Total
	Severe	Moderate	Mild	No ARDS	
Severe	56 (56.6)	0	0	0	56 (56.6)
Moderate	8 (8.1)	8 (8.1)	0	0	16 (16.2)
Mild	0	6 (6.1)	7 (7.1)	1 (1)	14 (14.1)
No ARDS	0	0	0	13 (13.1)	13 (13.1)
Total	64 (64.7)	14 (14.1)	7 (7.1)	14 (14.1)	99 (100)

Data are n (%). P/F_{1.0}=PaO₂/FiO₂ ratio with FiO₂ of 1.0; ARDS=Acute respiratory distress syndrome

DISCUSSION

The result of this study shows the P/F_{1.0} as an accurate measure for ARDS classification. Using P/F_{1.0} for ARDS classification yields an accuracy of about 85% compared to reference classification using pulmonary shunt fraction, while P/F ratios associated with FiO₂ ≤ 0.4 lead to accuracies lower than 40%. Indeed with increasing FiO₂ the rate of correct classification increases from <30% to about 85% which means that with increasing FiO₂ the calculated P/F ratio tends to be more correlated with the shunt fraction. Further experiments are required to study the effect of increasing inspiratory oxygen concentration on the accuracy of P/F ratio for classification of ARDS severity using experimental measurement of pulmonary shunt fraction.

The range of FiO₂ used in this study was wide enough to make obtaining a correlation with shunt level possible. Shunt fractions also showed a wide range suitable for the purpose of this study.

Shunt fraction estimated by OSA used in this study as the reference for ARDS severity classification. OSA has been used and validated extensively as a reliable tool for the estimation of gas exchange and acid-base parameters.^[15-25]

This study involved ARDS cases resulted from COVID-19. Previous studies had shown the effects of varying FiO₂ on the P/F ratios and on the resultant ARDS classifications in classic form of ARDS.^[8,10] The result of the present study shows the same effect for FiO₂ on P/F ratio and ARDS classification in COVID-19 pneumonia.

COVID-19 may involve the lungs and produces a wide spectrum of respiratory failure with limited therapeutic options at hand. Correct classification of COVID-19 induced ARDS severity based on P/F ratio may be important with respect to the provision of early invasive treatment options such as intubation, mechanical ventilation, prone position, and extracorporeal membrane oxygenation.^[26] We recommend using measured P/F_{1.0} for the classification of

ARDS severity. If the clinical measurement of P/F_{1.0} is not feasible, one may use the regression model presented in this study to estimate P/F_{1.0}. To determine the usefulness and the accuracy of the predictive model developed in this study for estimation of P/F_{1.0}, further clinical experiments on ARDS patients are necessary.

A possible limitation in the generalization of the result of this study is the fact that all enrolled cases were COVID-19 induced ARDS, while this may not significantly invalidate the study results, further study on non-COVID-19 cases and comparison with COVID-19 patients helps to shed more lights on the ARDS classification. Another concern is the fact that in addition many patients in this study were older than 60 years. Since the value of PaO₂ normally decreases with increasing age. Although the magnitude of this error is not large in patients under mechanical ventilation, anyhow this may lead to small errors when calculating P/F ratio.

CONCLUSION

ARDS severity classification using P/F_{1.0} shows the best match with the shunt-based classification. In addition, it is possible to use the regression model equation from this study to estimate the value of P/F_{1.0} when its clinical measurement is not practical.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Zetterström H. Assessment of the efficiency of pulmonary oxygenation. The choice of oxygenation index. *Acta Anaesthesiol Scand* 1988;32:579-84.
- Chang EM, Bretherick A, Drummond GB, Baillie JK. Predictive validity of a novel non-invasive estimation of effective shunt fraction in critically ill patients. *Intensive Care Med* 2019;7:49.
- Gowda MS, Klocke RA. Variability of indices of hypoxemia in adult respiratory distress syndrome. *Crit Care Med* 1997;25:41-5.
- Horovitz JH, Carrico CJ, Shires GT. Pulmonary response to major injury. *Arch Surg* 1974;108:349-55.
- Broccard AF. Making sense of the pressure of arterial oxygen to fractional inspired oxygen concentration ratio in patients with acute respiratory distress syndrome. *OA Crit Care* 2013;1:9.
- Lumb AB, Nunn JF. *Nunn's Applied Respiratory Physiology*. Amsterdam, The Netherlands: Elsevier; 2017. p. 246.
- ARDS Definition Task Force, Ranieri VM, Rubenfeld GD, Thompson BT, Ferguson ND, Caldwell E, et al. Acute respiratory distress syndrome: The Berlin Definition. *JAMA* 2012;307:2526-33.
- Karbing DS, Kjaergaard S, Smith BW, Espersen K, Allerød C, Andreassen S, et al. Variation in the PaO₂/FiO₂ ratio with FiO₂: Mathematical and experimental description, and clinical relevance. *Crit Care* 2007;11:R118.
- Siggaard-Andersen O, Siggaard-Andersen M. The oxygen status

- algorithm: A computer program for calculating and displaying pH and blood gas data. *Scand J Clin Lab Invest Suppl* 1990;203:29-45.
10. Pérez-Padilla R, Hernández-Cárdenas CM, Lugo-Goytia G. Classifying acute respiratory distress syndrome severity: Correcting the arterial oxygen partial pressure to fractional inspired oxygen at altitude. *Rev Invest Clin* 2016;68:169-70.
 11. Shirani K, Toghyani A. COVID-19 pneumonia with scant respiratory symptoms. *J Res Med Sci* 2020;25:82.
 12. Nokhodian Z, Ranjbar MM, Nasri P, Kassaian N, Shoaei P, Vakili B, *et al.* Current status of COVID-19 pandemic; characteristics, diagnosis, prevention, and treatment. *J Res Med Sci* 2020;25:101.
 13. Siggaard-Andersen M, Siggaard-Andersen O. Oxygen status algorithm, version 3, with some applications. *Acta Anaesthesiol Scand Suppl* 1995;107:13-20.
 14. Jibaja M, Ortiz-Ruiz G, García F, Garay-Fernández M, de Jesús Montelongo F, Martínez J, *et al.* Hospital mortality and effect of adjusting PaO₂/FiO₂ according to altitude above the sea level in acclimatized patients undergoing invasive mechanical ventilation. A multicenter study. *Arch Bronconeumol (Engl Ed)* 2020;56:218-24.
 15. Andreassen S, Egeberg J, Schröter MP, Andersen PT. Estimation of pulmonary diffusion resistance and shunt in an oxygen status model. *Comput Methods Programs Biomed* 1996;51:95-105
 16. Chu Z, Wang Y, You G, Wang Q, Ma N, Li B, *et al.* The P₅₀ value detected by the oxygenation-dissociation analyser and blood gas analyser. *Artif Cells Nanomed Biotechnol* 2020;48:867-74.
 17. Pérez-Padilla JR. Gas exchange at 2,240 m above sea level from computational models of the lung. *Neumol Cir Torax* 2018;77:19-23.
 18. Srinivasan AJ, Morkane C, Martin DS, Welsby IJ. Should modulation of p50 be a therapeutic target in the critically ill? *Expert Rev Hematol* 2017;10:449-58.
 19. Refsum HE, Opdahl H, Leraand S. Effect of extreme metabolic acidosis on oxygen delivery capacity of the blood – An *in vitro* investigation of changes in the oxyhemoglobin dissociation curve in blood with pH values of approximately 6.30. *Crit Care Med* 1997;25:1497-501.
 20. Nirmalan M, Willard T, Columb MO, Nightingale P. Effect of changes in arterial-mixed venous oxygen content difference (C (a-v) O₂) on indices of pulmonary oxygen transfer in a model ARDS lung. *Br J Anaesth* 2001;86:477-85.
 21. Mateják M, Kofránek J. “Physiomodel – An Integrative Physiology in Modelica,” 2015 37th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC), Milan, Italy; 2015. p. 1464-7.
 22. Dash RK, Korman B, Bassingthwaite JB. Simple accurate mathematical models of blood HbO₂ and HbCO₂ dissociation curves at varied physiological conditions: Evaluation and comparison with other models. *Eur J Appl Physiol* 2016;116:97-113.
 23. Wetterslev J, Hansen EG, Kamp-Jensen M, Roikjaer O, Kanstrup IL. PaO₂ during anaesthesia and years of smoking predict late postoperative hypoxaemia and complications after upper abdominal surgery in patients without preoperative cardiopulmonary dysfunction. *Acta Anaesthesiol Scand* 2000;44:9-16.
 24. Sowade O, Gross J, Sowade B, Warnke H, Franke W, Messinger D, *et al.* Evaluation of oxygen availability with oxygen status algorithm in patients undergoing open heart surgery treated with epoetin beta. *J Lab Clin Med* 1997;129:97-105.
 25. Siggaard-Andersen O, Ulrich A, Gøthgen IH. Classes of tissue hypoxia. *Acta Anaesthesiol Scand Suppl* 1995;107:137-42.
 26. Shaheen A, Tanaka D, Cavarocchi NC, Hirose H. Venovenous extracorporeal membrane oxygenation (VV ECMO): Indications, preprocedural considerations, and technique. *J Card Surg* 2016;31:248-52.