

# Effect of different dosages of nitroglycerin infusion on arterial blood gas tensions in patients undergoing on- pump coronary artery bypass graft surgery

Gholamreza Masoumi<sup>1</sup>, Evaz Hidar Pour<sup>2</sup>, Ali Sadeghpour<sup>3</sup>, Mohsen Ziaeyefard<sup>3</sup>, Mostapha Alavi<sup>3</sup>, Sanam Javid Anbardan<sup>4</sup>, Shahin shirani<sup>5</sup>

<sup>1</sup>Assistant Professor, Department of Cardiology, Shahid Rajaei Hospital, Isfahan University of Medical Science, Isfahan, Iran. <sup>2</sup>Professor, Department of Cardiology, Shahid Rajaei Hospital, Iran University of Medical Science, Tehran, Iran. <sup>3</sup>Assistant Professor, Department of Cardiology, Shahid Rajaei Hospital, Iran University of Medical Science, Tehran, Iran. <sup>4</sup>Student of Medicine, Tehran University of medical science, Tehran, Iran. <sup>5</sup>Assistant Professor, Department of Cardiology, Shahid Chamran Hospital, Isfahan University of Medical Science, Isfahan, Iran

**Background:** On-pump coronary artery bypass graft (CABG) surgery impairs gas exchange in the early postoperative period. The main object on this study was evaluation of changes in arterial blood gas values in patients underwent on pump CABG surgery receiving different dose of intravenous nitroglycerin (NTG). **Materials and Methods:** sixty-seven consecutive patients undergoing elective on-pump CABG randomly enrolled into three groups receiving NTG 50 µg/min (Group N1, n =67), 100 µg/min (Group N2, n = 67), and 150 µg/min (Group N3, n = 67). Arterial blood gas (ABG) tensions were evaluated just before induction of anesthesia, during anesthesia, at the end of warming up period, and 6 h after admission to the intensive care unit. **Results:** Pao<sub>2</sub> and PH had the highest value during surgery in Group N1, Group N2, and Group N3. No significant difference was noted in mean values of Pao<sub>2</sub> and PH during surgery between three groups ( $P > 0.05$ ). There was no significant difference in HCO<sub>3</sub> values in different time intervals among three groups ( $P > 0.05$ ). **Conclusion:** our results showed that infusing three different dosage of NTG (50, 100, and 150 µg/min) had no significant effect on ABG tensions in patients underwent on-pump CABG surgery.

**Key words:** Blood gas analysis, coronary artery bypass, nitroglycerin

## INTRODUCTION

Pulmonary function deterioration is the most frequent complication of on the pump coronary artery bypass graft (CABG) surgery in early postoperative phase.<sup>[1-4]</sup> It leads to impairment of gas exchange and prolonged extubation time.<sup>[5]</sup> On the other hand, respiratory dysfunction after cardiac surgery is the major cause of morbidity and mortality and increased duration of hospitalization.<sup>[6-8]</sup>

Occurrence of postoperative pulmonary problems is believed to be multifactorial. Some factors are patient dependent such as age, obesity, smoking<sup>[9]</sup> and others are surgical factors such as pain, general anesthesia, sternotomy, pleurotomy, medication, harvesting of the internal mammary artery, cardiopulmonary bypass, and alterations in lung and chest wall mechanics.<sup>[1,4,9-12]</sup>

It is thought that cardiopulmonary bypass induces many of these abnormalities by promoting an inflammatory response, increasing pulmonary endothelial permeability, and producing parenchymal damage.<sup>[13-16]</sup> Additionally, application of nitroglycerin – a potent vasodilator – in patients with coronary

artery disease, reduces PO<sub>2</sub> by increasing perfusion in hypoventilated areas and antagonizing hypoxic pulmonary vasoconstriction. Considering undesirable effect of CABG on the respiratory function (producing ventilation-perfusion mismatch, pulmonary shunting, pulmonary edema) and the frequent usage of nitroglycerin infusion in these patients that can decrease PaO<sub>2</sub> by pulmonary vasodilation and missing data regarding this issue, we designed this study to assess changes in arterial blood gas indices in patients receiving three different doses of intravenous nitroglycerin during on pump CABG surgery.

## MATERIAL AND METHODS

After obtaining institutional approval from Ethic Committee of our University and taking written informed consent from patients, this randomized double- blinded clinical trial was performed on 67 consecutive patients, aged between 50 and 75 years, who candidate for elective on pump CABG surgery. All enrolled patients had ejection fraction more than 40%, negative history for diabetes, renal, respiratory, and hematologic disorders. No patients had coexisting disease. If technique

**Address for correspondence:** Shahin Shirani, Assistant Professor, Department of Cardiology, Shahid Chamran Hospital, Isfahan University of Medical Science, Isfahan, Iran. E-mail: chancellor@mui.ac.ir

**Received:** 23.10.2011; **Revised:** 05.11.2011; **Accepted:** 15.11.2011

of anesthesia or surgery was changed, the patient was excluded from the study.

The study drugs were prepared with an anesthesiologist in similar syringe with similar volume. The patients had normal ABG and spirometry before surgery.

Patients were assigned randomly into three groups receiving NTG 50 µg/min (Group N1, n=67), NTG 100 µg/min (Group N2, n = 67), and NTG 150 µg/min (Group N3, n = 67).

Anesthesia was induced by midazolam (0.1 mg/kg), etomidate (0.2 mg/kg), sufentanil (1 µg/kg), and pancuronium (0.1 mg/kg) for all patients.

Blood for arterial gas analysis (PH, PaO<sub>2</sub>, PaCO<sub>2</sub>, HCO<sub>3</sub>), hemoglobin, and hematocrit was obtained in all patients at four points: just before induction of anesthesia, during anesthesia, at the end of warming up period and 6 h after admission to the intensive care unite (ICU).

Another anesthesiologist who was not aware from the study drug preparation recorded the data.

It was calculated that sample size of 67 patients in each

group would have an 80% power to detect a 24% difference in the mean partial pressure of oxygen in the blood among three groups, with Alfa=0.05. Differences among three groups mean were compared using one-way analysis of variance (ANOVA) and posthoc comparisons at various points in time by using Bonferroni's type I error rate correction for multiple tests of significance. Categorical variables were analyzed by Pearson chi-square test and by Fisher's exact test when the anticipated number was less than 5. P less than 0.05 was considered statistically significant. All statistical analyses were performed using SPSS 16.0 for Windows statistical package.

## RESULTS

In our clinical trial, 67 patients were evaluated (mean age: 60.2 ± 10.29). Forty-one patients (61.2%) were male and 26 (38.8%) were female. All patients underwent on pump CABG surgery and the internal mammary artery was used for all. The flow diagram of randomized patients is shown in [Figure 1].

The mean duration of surgery was 3.4 ± 0.8 h. As mentioned in inclusion criteria, all patients had ejection fraction more than 40% and had no coexisting disease.




Figure 1: Flow diagram of randomized patients. NTG = nitroglycerin


Regarding to our results,  $Po_2$  and PH had the highest value during surgery and in preoperative phase in all three groups, respectively. On the other hand,  $HCO_3$  demonstrated subtle changes during all phases with no significant changes among three groups. Results are demonstrated [Figures 1–6].

## DISCUSSION


Patients undergoing elective on-pump CABG enrolled in this study. Arterial blood gas status was evaluated in four different points in order to assess influence of different doses of intravenous nitroglycerin.

Patients' hemoglobin decreased during operation and the mean hemoglobin was lowest during operation ( $Hb: 7.09 \pm 0.09$  mg/dL) and hemoglobin changes were statistically significant during different phases of study with no significant difference among different groups. Furthermore, the highest hematocrit was detected in preoperative phase and its changes were statistically significant during different phases of study but not significant among different groups. It seems that blood loss during surgery is responsible for this phenomenon.


On the other hand,  $PO_2$  was maximum during operation and decreased during warming up period and ICU admission but did not reached lower levels than first values.




**Figure 1:** PH During different phases with different doses of intravenous nitroglycerin, NTG = nitroglycerin; ICU = intensive care unit; The unit of NTG dosage is microgram; pH = arterial pH




**Figure 2:** Po2 in different phases regarding different doses of NTG, NTG = nitroglycerin; ICU = intensive care unit; The unit of NTG dosage is microgram; Po2 = arterial Po2




**Figure 3:** PCO2 during different phases with different doses of intravenous nitroglycerin, NTG = nitroglycerin; ICU = intensive care unit; The unit of NTG dosage is microgram; Pco2 = arterial Pco2



**Figure 4:** HCO3 during different phases with different doses of intravenous nitroglycerin, NTG = nitroglycerin; ICU = intensive care unit; The unit of NTG dosage is microgram; HCO3 = arterial HCO



**Figure 5:** Hemoglobin (Hb) during different phases with different doses of intravenous nitroglycerin, NTG = nitroglycerin; ICU = intensive care unit; The unit of NTG dosage is microgram



**Figure 6:** Hematocrit during different phases with different doses of intravenous nitroglycerin, NTG = nitroglycerin; ICU = intensive care unit; The unit of NTG dosage is microgram

Additionally, the mean arterial PO<sub>2</sub> was more than 90 mmHg in all groups, during all phases and no significant difference was found. It seems that hypoxemia did not take place during operation and early postoperative phase (first 6 h).

Considering our results, it seems that no prominent changes in ABG occur during the first few hour – 6 h in our study – after CABG, while previous studies, reported reduction in arterial PO<sub>2</sub> common following cardiac surgeries that persists for few weeks after surgery and is lowest on second day after operation then improves gradually.<sup>[17-23]</sup>

Masoumi *et al.*<sup>[24]</sup> in a clinical trial showed that applying different doses of intravenous NTG did not exert a significant influence on cardiac instability and the need for use of inotrope drugs. Their conclusion can be one of causes of our finding.

Singh and colleagues<sup>[9]</sup> showed that after CABG surgery, there is a marked decrease in PaO<sub>2</sub>. The decrease in PaO<sub>2</sub> occurs usually on the second postoperative day and improves subsequently but remains below preoperative values on the eighth day postoperatively.

The decrease in PaO<sub>2</sub> can be due to decrease in hematocrit and increased permeability pulmonary edema that noted in the postoperative period.<sup>[19]</sup>

Our study has two limitation: firstly, lack of data about the alveolar arterial gradient – an important factor for evaluating the postoperative respiratory function – may contribute to our results while in similar studies it is considered as a mainstay for determination of patient's ventilation system.

Secondly, 6 h is a short duration for evaluating patients for changes in ABG and hypoxemia so more studies is recommended with longer follow up time and more accurate measurements.

## ACKNOWLEDGMENT

The authors would like to thank Farzan institute for their support in performing this study.

## REFERENCES

1. Taggart DP. Respiratory dysfunction after cardiac surgery: Effects of avoiding cardiopulmonary bypass and the use of bilateral internal mammary arteries. *Eur J Cardiothoracic Surg* 2000;18:31-7.
2. Lichtenberg A, Hagl C, Harringer W, Klima U, Haverich A. Effects of minimal invasive coronary artery bypass on pulmonary function and postoperative pain. *Ann Thorac Surg* 2000;70:461-5.
3. Berrizbeitia LD, Tessler S, Jacobowitz IJ, Kaplan P, Budzilowicz L, Cunningham JN. Effect of sternotomy and coronary bypass surgery on postoperative pulmonary mechanics. *Chest* 1989;96:873-6.
4. Cimen S, Ozkul V, Ketenci B, Yurtseven N, Günay R, Ketenci B, *et al.* Daily comparison of respiratory functions between on-pump and off-pump patients undergoing CABG. *Eur J Cardiothorac Surg* 2003;23:589-94.
5. Montes FR, Maldonado JD, Paez S, Ariza F. Off-Pump versus on-pump coronary artery bypass surgery and postoperative pulmonary dysfunction. *J Cardiothorac Vasc Anesth* 2004; 18:698-703.
6. Messent M, Sullivan K, Keogh BF, Morgan CJ, Evans TW. Adult respiratory distress syndrome following cardiopulmonary bypass: Incidence and prediction. *Anaesthesia* 1992;47:267-8.
7. Hachenberg T, Tenling A, Hansson HE, Tydén H, Hedenstierna G. The ventilation-perfusion relation and gas exchange in mitral valve disease and coronary artery disease: Implications for anesthesia, extracorporeal circulation, and cardiac surgery. *Anesthesiology* 1997;86:809-17.
8. Hulzebos EH, Helders PJ, Favié NJ, De Bie RA, Brutel de la Riviere A, Van Meeteren NL. Preoperative intensive inspiratory muscle training to prevent postoperative pulmonary complications in high-risk patients undergoing CABG surgery: A randomized clinical trial. *JAMA* 2006;296:1851-7.
9. Arabaci U, Akdur H, Yiğit Z. Effects of smoking on pulmonary functions and arterial blood gases following coronary artery surgery in Turkish patients. *Jpn Heart J* 2003;44:61-72.
10. Rolla G, Fogliati P, Bucca C, Brussino L, Di Rosa E, Di Summa M, *et al.* Effect of pleurotomy on pulmonary function after coronary artery bypass grafting with internal mammary artery. *Respir Med* 1994;88:417-20.
11. Vargas FS, Terra-Filho M, Hueb W, Teixeira LR, Cukier A, Light RW, *et al.* Pulmonary function after coronary artery bypass surgery. *Respir Med* 1997;91:629-33.
12. Hachenberg T, Tenling A, Nystrom SO, Tyden H, Hedenstierna G. Ventilation-perfusion inequality in patients undergoing cardiac surgery. *Anesthesiology* 1994;80:509-19.
13. Wan S, LeClerc JL, Vincent JL. Inflammatory response to cardiopulmonary bypass: Mechanisms involved and possible therapeutic strategies. *Chest* 1997;112:676-92.
14. Hall RI, Smith MS, Rucker G. The systemic inflammatory response to cardiopulmonary bypass: Pathophysiological, therapeutic, and pharmacological considerations. *Anesth Analg* 1997;85:766-82.
15. Paparella D, Yau TM, Young E. Cardiopulmonary bypass induced inflammation: Pathophysiology and treatment; an update. *Eur J Cardiothorac Surg* 2002;21:232-44.
16. Miura N, Yoshitani K, Kawaguchi M, Shinzawa M, Irie T, Uchida O, *et al.* Jugular bulb desaturation during off-pump coronary artery bypass surgery. *J Anesth* 2009;23:477-82.
17. Quinlan GJ, Mumby S, Lamb NJ, Moran LK, Evans TW, Gutteridge JM. Acute respiratory distress syndrome secondary to cardiopulmonary bypass: Do compromised plasma iron-binding anti-oxidant protection and thiol levels influence outcome? *Crit Care Med* 2000;28:2271-6.
18. Braun SR, Birnbaum ML, Chopra PS. Pre and post-operative pulmonary function abnormalities in coronary artery revascularization surgery. *Chest* 1978;73:316-20.
19. Singh NP, Vargas FS, Cukier A, Terra-Filho M, Teixeira LR, Light RW. Arterial blood gases after coronary artery bypass surgery. *Chest* 1992;102:1337-41.
20. Lonsky V, Svitek V, Brzek V, Kubicek J, Volt M, Horak M, Mandak J. Direct oxymetric peripheral tissue perfusion monitoring during open heart surgery with the use of cardiopulmonary bypass: Preliminary experience. *Perfusion* 2011;26:510-5.
21. Moritz S, Rochon J, Völkel S, Hilker M, Hobbhahn J, Graf BM, *et al.* Determinants of cerebral oximetry in patients undergoing off-pump coronary artery bypass grafting: An observational study.

- Eur J Anaesthesiol 2010;27:542-9.
22. Javidi D, Saffarian N. Prognostic value of left ventricular myocardial performance index in patients undergoing coronary artery bypass graft surgery. Arch Iran Med 2008;11:497- 501.
  23. Taghipour HR, Naseri MH, Safiarian R, Dadjoo Y, Pishgoo B, Mohebbi HA, *et al.* Quality of life one year after coronary artery bypass graft surgery. Iran Red Crescent Med J 2011;13:171-7.
  24. Masoumi G, Hidarpour E, Tabae AS, Ziaeyefard M, Azarasa

A, Abneshahidi A, *et al.* Evaluating hemodynamic outcomes of different dosages of intravenous nitroglycerin after coronary artery bypass graft surgery. J Res Med Sci 2011;16:910-5.

**How to cite this article:** Masoumi Gh, Hidar Pour E, Sadeghpour A, Ziaeyefard M, Alavi M, Javid Anbardan S *et al.* Effect of different dosages of nitroglycerin infusion on arterial blood gas tensions in patients undergoing on-pump coronary artery bypass graft surgery. J Res Med Sci 2012; 17(2): 123-7.

**Source of Support:** Nil, **Conflict of Interest:** None declared.