

Response rates to HB vaccine in CKD stages 3-4 and hemodialysis patients

Mohammad H. Ghadiani, Shahin Besharati², Nouraddin Mousavinasab³, Mojgan Jalalzadeh¹

Department of Nephrology, Taleghani Hospital, Departments of Nephrology, Imam Hossein Hospital, Shahid Beheshti University of Medical Sciences, Tehran, ²Department of Internal Medicine, Vali-e-asr Hospital, Zanjan University of Medical Sciences, ³Department of Social Medicine, School Medicine, Zanjan University of Medical Sciences, Zanjan, Iran

Background: Hepatitis-B virus (HBV) infection is a big problem in chronic kidney disease (CKD) population. We attempted to compare the response rate to HB vaccine in CKD stages 3-4 patients with that in hemodialysis (CKD stage-5; HD patients) and medical staff. **Materials and Methods:** Three hundred and three participants were enrolled into the study to test the seroconversion rate after vaccination. Participants formed three groups: Group-A: HD patients, Group-B: diagnosed with CKD stages 3-4, and Group-C: healthy medical staff. CKD stages 3-4 participants were vaccinated from February to November 2010. HD patients were vaccinated at the time of initial HD. While the medical staffs were vaccinated at the time they started working at the hospital. Group-A, Group-B and Group-C received four 40 μ g (in 0,1,2 and 6 months), three 40 μ g (0, 1 and 6 months) and three 20 μ g (0, 1 and 6 months) doses of HB vaccine, respectively. Three months after completion of the vaccination schedule, seroconversion and seroprotection rates in each group were investigated. **Results:** Seroconversion rates were 44.3%, 89.7%, and 96.2% for groups A, B and C, respectively. CKD stages 3-4 patients showed higher response rate than dialysis patients [$\chi^2(1):30.6, P < 0.001$]. But a significant difference in the seroconversion rate between CKD stages 3-4 patients and medical staffs was not observed [$\chi^2(1):3.4, P = 0.064$]. Multivariate analyses showed patients with more advanced CKD and who were older had less seroconversion rates [odds ratio: 0.09(95%CI: 0.04 - 0.25) and [odds ratio: 0.39(95% CI: 0.18-0.85)], respectively. But sex was not associated with seroconversion ($P > 0.05$). **Conclusion:** Stages 3-4 patients with higher dosages of routine HB vaccine had higher seroconversion rate than HD patients. Future studies should evaluate the recommended dosage of HB vaccine among these patients.

Key words: Chronic kidney disease, hemodialysis, Hepatitis-B vaccine, seroconversion

INTRODUCTION

One of the most serious infectious diseases in the world is Hepatitis-B virus (HBV). It is estimated that 200–500 million people are infected by the virus in the world;^[1] 360 million have chronic infection and 600,000 die by acute hepatitis, liver fibrosis, or hepatocellular carcinoma each year.^[2] Hepatitis-B infection is difficult to treat, that is why prevention by Hepatitis-B vaccine (HB vaccine) is the most efficient way to tackle the problem. HB vaccine has been available since 1982.^[3] Today's HB vaccine contains 20 μ g HB surface antigen (HBsAg) along with 0.5 mg aluminum salt as adjuvant. In the general population, seroprotection is defined by an HB surface antibody (HBsAb) titer greater than 10 mIU/ml. The HB vaccine has been confirmed to be

safe, and there is no evidence indicating it causes any complications or side effects. Seroprotection rates have been attained in up to 95% of healthy individuals with a 23 standard dose of vaccination.^[2] In another study on Iranian population standard HBV vaccination has resulted in an immune response in 57.9% of subjects.^[4] There are some studies about the prevalence of chronic HBV infection in general population of Iran.^[5,6] These studies have shown that because of the vaccination of newborns against HBV, since 1992, prevalence of chronic HBV infection has reduced.

Due to exposure to blood products, hemodialysis (HD) patients have a higher risk for several infections.^[7] These patients are at increased risk for becoming carriers of the HBV, which would lead to eventual cirrhosis and liver carcinoma.^[8]

Because of erythropoietin, the incidence of HB infection has reduced.^[9] However, occurrence of HB infection in HD units continues to occur with different frequencies.^[7,10-14] HB vaccine has reduced the number of HB infections in patients with chronic kidney disease (CKD).^[15]

Access this article online	
Quick Response Code:	Website:
	www.journals.mui.ac.ir/jrms
	DOI:

Address for correspondence: Dr. Mojgan Jalalzadeh, Assistant Professor, Department of Nephrology and Dialysis, Imam Hossein Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran. E-mail: j_mojgan@yahoo.com

Received: 19-05-2012; **Revised:** 12-06-2012; **Accepted:** 14-06-2012

In HD patients, the effectiveness of HB vaccine to seroconversion is lower than in the general population.^[16-19] Patients with end-stage renal disease (ESRD) have several defects in both the humoral and cellular immune response,^[20,21] which results in reduced production of antibody.^[7,15,22]

One study has suggested that to improve this response rate, patients should be vaccinated at the early stages of CKD and before requiring dialysis or transplantation.^[23]

In one Iranian population study, the response rate to vaccination in predialysis patients was the same as in dialysis patients.^[24]

One of the largest studies performed on this issue enrolled 61 renal failure patients. The authors concluded that patients who were not on dialysis had a better immune response to HB vaccine than the patients who were on dialysis.^[25] However, the best strategy for vaccination at CKD stages 3-4 patients before HD has not been ascertained yet.

Our current prospective study aimed to compare seroconversion rates between three groups of patients namely, CKD stages 3-4, under HD and medical staff as the control group. We also hoped to be able to identify some of the possible parameters that could influence the response rates to HB vaccine.

MATERIALS AND METHODS

In this clinical trial 303 participants have been involved. According to the glomerular filtration rate (GFR), the patients were divided into three groups: HD patients (Group-A, GFR < 15 ml/min), patients with CKD stages 3-4 (Group-B, 15 < GFR < 60 ml/min), and medical staff (Group-C, GFR > 90 ml/min). All CKD and HD patients were receiving care at a kidney dialysis center in the Iranian provincial capital of Zanjaan.

All CKD participants were vaccinated from February to November 2010. HD patients were vaccinated at the time of initial HD and medical staff at the time of being hired.

The vaccination protocol for groups A and C was as follows: Group-A received 40 µg at 0, 1, 2 and 6 months and Group-C received 20 µg at 0, 1 and 6 months. In addition, the vaccines were injected into the deltoid muscle and according to the policy of the hospitals and HD wards the levels of antibody production of both groups were checked 3 months after the last dose of injection.

We used both the Cockcroft-Gault (C-G) and modified diet in renal disease (MDRD) formulas to determine kidney function, GFR.

For Group-B, the entry criteria included a GFR of less than 60 ml/min and more than 15 ml/min, as was determined by using the C-G formula $[(140 - \text{age}) \times \text{lean body weight} / \text{serum creatinine} \times 72]$ and estimated glomerular filtration rate (eGFR), which was calculated by using the abbreviated modified diet in renal disease (MDRD) formula, where $\text{eGFR (ml/min)} / 1.73 \text{m}^2 = 186 \times (\text{Creatinine}/88.4)^{-1.154} \times (\text{age})^{-0.203} \times 0.742$ (if female) or 1.21 (if African American).^[15] Patients in Group-B were withdrawn from the study if their renal function had deteriorated to the point that dialysis treatment was needed.

The exclusion criteria for Group-B included: hypersensitivity to HB vaccine, a history of previous HB infection and taking immune suppressive medications as well as patients with known lymphoproliferative disorder.

All participants in Group-B signed a consent form after being informed about the study. Before enrollment, patients were tested for HBsAg, anti-HBc and anti-HBs antibodies by enzyme-linked immunosorbent assay (ELISA) methods. All patients of Group-B who were identified as HB infection negative and anti-HBs negative started a vaccination schedule consisting of recombinant vaccine 40 µg at 1, 2 and 6 months. Vaccine was administered intramuscularly into the deltoid muscle. The vaccine was well tolerated; no patient had a serious adverse event attributable to the vaccine, and no patient withdrew from the study because of an adverse event. All patients received their vaccinations at the clinic.

HBsAb was measured by using ELISA methods at virology laboratory of Vali-e-Asr Hospital. Anti-HBs titers are expressed in mIU/ml. For all the participants, Anti-HBs antibodies were measured 3 months after the completion of the vaccine schedule. Anti-HBs titers less than 10 mIU/ml were defined as non-seroconversion. Antibody titers more than 10 mIU/ml were considered as seroconversion, and HBs titers above 100 mIU/ml were considered seroprotective level, for groups A and B. Antibody titers more than 10 mIU/ml were considered as seroprotection for Group-C. For all groups, patients with antibody levels < 10 mIU/ml were considered non-responders.

All the HD patients were on bicarbonate HD for 3.5 h, three times a week, by using 1.72.1 m² capillary high flux polysulfone dialyzer and a dialysate flow rate of 300 and 350 ml/min. Hematocrit levels were maintained between 30 and 38% by administration of appropriate doses of recombinant erythropoietin.

The study was approved by the research ethics committee of the Zanjaan University of Medical Science (Ethics No: 19/3-3/236). Trial Registration Number: IRCT 138905163325N4

Statistical analysis

Baseline characteristics are presented as mean \pm SD or median with 25th and 75th percentiles, depending on the underlying distribution. Continuous variables were compared in three groups using analysis of variance or KruskalWallis test, when appropriate, while discrete variables were compared by using chi-square test. We used *t*-test or Mann–Whitney U-test (for non-parametric samples) and chi-square test to compare baseline characteristics between seroconverters and non-seroconverters. A *P*-value of less than 0.05 for two-sided univariate test was considered statistically significant.

A logistic regression model was used to identify important predictors of seroconversion. Variables included in multivariate analysis were patients' age, sex and GFR. All models were tested for interaction.

RESULTS

Three hundred three HBsAg negative participants [101 males (33.3%) and 202 females (66.6%), with a mean age of 44.30 years] were enrolled for this study. Table-1 presents the characteristics of the study population.

Table 1: Characteristics of study population at enrollment

Variables	No. (%)
Age (years)	44.3 \pm 17.9 (30-59)*
Age > 60 years	73 (24.1)
Male gender	101 (33.3)
Weight (kg)	68.7 \pm 10.9 (60-78)
Creatinine (mg/dl)	2.3 \pm 1.9 (0.8-3.8)
GFR (CockcroftGault) (ml/min/1.73 m ²)	69.4 \pm 48.8 (14.6-108.3)
GFR (MDRD) (ml/min/1.73 m ²)	57.0 \pm 39.7 (15.2-90.1)
Anti-HBs levels (mIU/ml)	224.4 \pm 259.1 (20-324)
Seroconversion rate	242 (79.9)
Seroprotection rate	206 (68)

*Values expressed as Means SD (25th-75th percentiles) and numbers (percentage)

Table 2 lists and compares demographic and laboratory data of the three study groups. Group-A comprised 49 males (55.7%) and 39(44.3%) females with a mean age of 58.8 years. Group-B consisted of 29 males (50%) and 29 females (50%), with a mean age of 57.7 years. Group-C included 23 males (14.6%) and 134 females (85.4%), with a mean age of 31.2 years.

Mean serum creatinine level of CKD patients (Group-B) was 2.12 \pm 0.61 mg/dl; according to C-G formula, the creatinine clearance amounted to a GFR mean of 35.4 \pm 12 ml/min with an eGFR of 32.2 \pm 9.9 ml/min/1.73 m² as calculated by MDRD formula. Based on the above data, the participants of Group-B had stage 3 and/or stage 4 of CKD.

For HD patients (Group-A) the GFR mean according C-G formula was 13.4 \pm 1.3 ml/min, while the GFR mean according MDRD formula was 11.9 \pm 3.3 ml/min/1.73 m².

GFR according both MDRD and C-G was lower in HD patients [*P* < 0.001, 95% confidence interval (CI): 19-24]. There were no significant gender and age differences between groups A and B.

Compared to groups A and B, participants in Group-C were younger [*P* < 0.001, *t* (243): 20.5 and *t* (213):17.2], and there were more females [χ^2 (1):45.73 and 28.8, *P* < 0.001] than males.

Two hundred and forty-two individuals out of the 303 participants in the study (79.9%) seroconverted after the vaccine schedule (anti-HBs levels \geq 10 mIU/ml) as measured 3 months after the last vaccine was administered. The mean peak of anti-HBs titer was 224.4 \pm 259.1 IU/l (20324 IU/l) [Table 1].

Seroconversion rate in HD patients was 44.3%, for Group-B patients 89.7% [χ^2 (1):30.6, *P* < 0.001] and for Group-C 96.2% [χ^2 (1):87.1, *P* < 0.001]. A significant difference in seroconversion rate between Group-B patients and medical staff was not observed (χ^2 (1):3.4, *P* < 0.064) [Table 2].

Table 2: Comparison of different variables between study groups

Variables	Group-A (HD patients) (n = 88) (%)	Group-B (CKD patients) (n = 58) (%)	Group-C (Medical staffs) (n = 157) (%)	<i>P</i> value
Age (years)	58.8 \pm 15.4	57.7 \pm 17.2	31.2 \pm 5.2	<0.001 F (2,300): 200.9
Age > 60 years	49 (55.7)	24 (41.4)	0	<0.001 χ^2 (2):107.3 ^a
Male gender	49 (55.7)	29 (50)	23 (14.6)	<0.001 χ^2 (2):51.7
Weight (kg)	64.6 \pm 11	68.0 \pm 11.0	71.2 \pm 10.1	<0.001 F (2,300): 11.1
Creatinine (mg/dl)	9.05 \pm 2.3	2.12 \pm 0.6	0.84 \pm 0.17	<0.001 F (2,300): 850
GFR (Cauckrof-Gault) (mL/min/1.73 m ²)	13.5 \pm 1.3	35.4 \pm 12	113.3 \pm 20.7	<0.001 F (2,300): 1283
GFR (MDRD) (ml/min/1.73 m ²)	11.9 \pm 3.3	32.1 \pm 9.9	91.5 \pm 21.1	<0.001 F (2,300): 789.2
Anti-HBs levels (mIU/ml)	7.95 (5.1-161)	123 (33.1-540.3)	242 (61.5-313.5)	<0.001 χ^2 (2):47.9
Seroconversion rate	39 (44.3)	52 (89.7)	151 (96.2)	<0.001 χ^2 (2):95.6
Seroprotection rate	23 (26.1)	32 (55.2)	151 (96.2)	<0.001 χ^2 (2):132.5

HD = Hemodialysis, CKD = Chronic kidney disease, GFR = Glomerular filtration rate; Values are expressed as Mean \pm SD for continuous variables with normal distribution and as median rank (25th-75th percentiles) for continuous variables without normal distribution. Categorical variables are expressed as number(percentage) Chi-square value(df)

Seroconversion (anti-HBs levels > 100 mIU/ml) occurred in 55.2% of Group-B patients (32 out of 58) and in 26.1% of HD patients (23 out of 88) and 96.2% of Group-C (151 out of 157) ($P < 0.001$).

Univariate analysis

Table 3 shows the comparison data of seroconverters and nonseroconverters.

Out of the 303 vaccinated participants, 44.3% of HD (39 patients), 89.7% of CKD stages 3-4 (52 patients) and 96.2% (151) of participants in the medical staff group were seroconverted [Table 2].

Seroconversion and renal function

When GFR decreased, there was a clear statistically significant decline in seroconversion. This was the case regardless of whether the C-G (not shown) or MDRD formula was used.

Univariate analysis also indicated that the severity of renal function had a significant effect on seroconversion rates. The seroconversion rates for patients with poor renal function were significantly different from those with better renal function, whether calculated by C-G {creatinine clearance <15 ml/min, seroconversion = 39/88, 44%; creatinine clearance = 15-60 ml/min, seroconversion = 52/58, 90% and creatinine clearance >90 ml/min, seroconversion = 151/157, 96% [$\chi^2(2) = 95.6, P < 0.001$] or by MDRD {(eGFR <15 ml/min/1.73 m² 35/73, 48%; eGFR 15-60 ml/min/1.73 m² 58/76, 76%; and eGFR >90 ml/min/1.73 m² 76/77, 99% [$\chi^2(2) = 66.4, P < 0.001$]}.

Table 3 also shows that patients with higher GFR levels had significantly more seroconverters. Univariate analysis also showed that subjects with GFR>90 were 31 times more likely to have seroconversion than subjects with GFR<15 (95%CI: 12.776.9, χ^2 :54.3, $P < 0.001$) and the odds of seroconversion for subjects with GFR>90 were three

times higher than for those with GFR=15-60 (95%CI: 0.99, χ^2 :3.2, $P < 0.08$).

Seroconversion and age

The mean age of the seroconverter group (40.6 ± 16 years) was lower than that of the non-seroconverter group (59.0 ± 17 years). This difference was significant when univariate analysis was used [$t(301)$: 7.8, $P < 0.001$].

Fifty-nine percent of patients in non-seroconverter group were above 60 years compared with 15% of seroconverters [Table 3].

The univariate analysis indicated that individuals under 60 years of age had significantly higher seroconversion (89%) than those above 60 years (51%) [$\chi^2(1)$:43.2, $P < 0.001$].

Those under 60 years were eight times more likely to be seroconverters (95%CI: 4.3-14.8).

Seroconversion and gender

Fifty-three percent of patients in the non-seroconverter group were males, while 29% of seroconverters were males (χ^2 :12.6, $P < 0.001$).

Univariate analysis indicated that females had significantly higher seroconversion (85.6%) than males (68.3%) (χ^2 :12.0, $P = 0.001$). Females had 2.8 times higher odds for seroconversion compared to males (95%CI: 1.6-4.9).

Multivariable analysis

Logistic regression analysis was used to predict the probability of seroconversion and factors that influenced such a conversion. The predictor variables were the participants' gender, age and three GFR categories.

The model was able to correctly classify 92.6% of those who seroconverted and 50.8% of those who did not, with an overall success rate of 84.2%.

Table 3: Univariate comparison by seroconversion

Variables	Non-seroconversion (n = 61) (%)	Seroconversion (n = 242) (%)	P value
Age (years)	59.0 ± 17.5	40.6 ± 16.2	<0.001 t (301):7.8
Age > 60 years	36 (59.0)	37 (15.3)	<0.001 $\chi^2(1)$:50.9
Male gender	32 (52.5)	69 (28.5)	<0.001 $\chi^2(1)$:12.6
Weight (kg)	66.4 ± 11.9	69.3 ± 10.6	0.062 t (301): -1.8
Creatinine (mg/dl)	4.3 ± 1.8	1.8 ± 1.7	<0.001t (301): 10.1
GFR (CockcroftGault) (ml/min/1.73 m ²)	14.07 (13.0-14.8)	96.52 (31.8-111.6)	<0.001 Z: 8.9
GFR (MDRD) (ml/min/1.73 m ²)	12.73 (9.9-19.7)	72.68 (31.0-92.3)	<0.001 Z: 8.4
GFR<15 cc/min	49 (80.3)	39 (16.1)	<0.001 $\chi^2(1)$:97.5
GFR>90 cc/min	6 (9.8)	151 (62.4)	<0.001 $\chi^2(1)$:53.9
15≤GFR<60 cc/min	6 (9.8)	52 (21.5)	0.039 $\chi^2(1)$:4.2

HD = Hemodialysis, CKD = Chronic kidney disease, GFR = Glomerular filtration rate; Values are expressed as Mean±SD for continuous variables with normal distribution and as median (25th75th percentiles) for continuous variables without normal distribution. Categorical variables are expressed as number(percent) Chi-square value(df)

Table 4: Multivariate modeling results for predictors of seroconversion

Variables	Odds ratio	95%CI	Wald (χ^2)	P value
Sex (male)	0.88	0.42-1.8	0.11	0.74
Age > 60 years	0.38	0.18-0.84	5.6	0.017
GFR >90cc/min			44.7	<0.001
GFR<15-60 cc/min	0.056	0.2-0.16	28.82	<0.001
GFR <15 cc/min	0.583	0.15-2.1	0.65	0.41
Constant	1.42		59.4	<0.001

Table 4 shows the logistic regression coefficients, Wald test and odds ratio with 95% CI for each of the predictors.

Using a 0.05 criterion of statistical significance, age>60 years and GFR <15 (HD group) had significant partial effects.

Those with the lowest level of kidney function were less likely to seroconvert than those with better kidney function, independent of other factors.

The significant odds ratios for older age, when holding all other variables constant, were found. Inverting the odds ratio for age variable revealed that the under 60-year-old participants were 2.6 times more likely to seroconvert than individuals above 60 years of age (95%CI: 1.2-5.6, χ^2 :5.68, $P=0.017$). Male sex was adjusted for, but was not significant in the model.

Logistic regression with adjusted gender and age showed that the chance for development of seroconversion for the patients with 15<GFR <60 decreased to about 0.95 compared to the subjects with GFR>90. This is while, the rate of development for seroconversion for those with GFR <15 declined to 0.42 compared with subjects with GFR>90.

In Table 4, gender did not show any effect on producing seroconversion.

DISCUSSION

HepatitisB vaccine should be administered to all ESRD patients and healthcare workers in this unit.^[26,27] Due to impaired immune system, patients with CKD have a less than optimal response to HB vaccine.^[26] The recommended vaccination schedule in dialysis patients is four doses of 40 μ g, which is twice the normal dosage, at intervals of 0, 1, 2, and 6 months to complete the primary immunization series.^[20] To increase the effectiveness of HB vaccine, various methods have been tested, such as recombinant adjuvants, thymopentine, levamisole, zinc, interleukin-2, interferon, erythropoietin, granulocyte-macrophage colony-stimulating factor (GM-CSF), reinforced intramuscular(IM)/intra dermal(ID) methods and new vaccines that contain adjuvant, all of which function differently.^[28-36]

At present, it is recommended that anti-HBs antibody titers be kept above 100 mIU/ml to induce a protective antibody response in HD patients.^[34] Patients who develop anti-HBs \geq 100 mIU/ml to HB vaccine should be given a booster dose of vaccine every 5 years. Patients with low level antibody response (anti-HBs \geq 10 mIU/ml or <100 mIU/ml) should be given a booster shot after 1 year and every 5 years thereafter.^[37]

Until now, there have been no clear data available on the vaccination of patients with CKD stages 3-4, who have a 15 < GFR < 60 ml/min. As it can be seen in the subsequent studies, various dosages, frequencies and methods of administration of HB vaccine in this population have been utilized to develop a protective antibody response.

This study was conducted to determine the efficacy of three doses of 40 μ g vaccine in stages 3-4 of CKD patients at the time of diagnoses, as compared with the threedoses of 20 μ g vaccine schedule given to the normal healthy medical staff and fourdoses of 40 μ g vaccine schedule given to the HD population. Since responsiveness to HB vaccine in patients with renal disease is reduced, in order to assess a better response, higher dosage of HB vaccine (40 μ g) was administered.

C. A. M. McNulty and his colleagues compared 20 and 40 μ g doses of HB vaccine, administered at 0, 1 and 6 months to 121 patients with moderate renal insufficiency. The subjects were at stages 2 and 3 of CKD. Seroconversion was greater (67%) after three doses of 40 μ g of vaccine than in patients given a standard dose of 20 μ g (57%). However, this was not statistically significant.^[37] They showed there was no correlation between the seroconversion rate and severity of CKD.

Gerald DaRoza *et al.* studied 165 patients with GFR<30 ml/min. They vaccinated patients with 40 μ g IM at 0, 1, and 6 months. They found an association between the GFR and the seroconversion rate in the subjects.^[19]

In our study, seroconversion rates in patients with GFR<15, GFR1560 and GFR>90 ml/min were 44.3%, 89.7%, and 96.2%, respectively. This study showed that the degree of renal impairment has an influence on seroconversion rate.

Siddiqui and colleagues studied the efficacy of four doses of 40 µg vaccine in 130 CKD patients as compared to the three-dose 20 µg schedule for the normal population. They found that administering 40 µg vaccine produced a better seroprotection in CKD patients.^[38]

Our current study showed 96.2%, 55.2% and 26.1% seroprotection in medical staff, stages 3-4 of CKD and HD groups, respectively.

Fraser *et al.* conducted a study of 68 patients with different severities of renal failure. Patients received four doses of HB vaccine, after which anti-HBs titers were checked at 0, 1, 2, 3, 6, 8 and 12 months. Maximum anti-HBs titers were observed at 8 months. They reported that the degree of renal failure had an effect on the titer of anti-HBs ($P < 0.002$).^[39]

In our study, patients with maximum renal failure, those being the HD patients, had less seroconversion ($P < 0.002$) and seroprotection ($P < 0.002$).

In another study, 78 children and adolescents with CKD were given three 20-µg dose course of the HB vaccine.^[40] After only two doses of vaccine, seroconversion appeared in all patients. The results showed that a regimen of three 20-µg doses of HB vaccine produced proper immunogenic response in children with CKD. The researchers recommended using three doses of vaccine before the development of ESRD.^[40]

Some factors could possibly influence the antibody levels in CKD patients. Older age, male sex, uremia inadequate dialysis, nutritional status, body weight, presence of diabetes, use of low biocompatibility dialysis material, hyperparathyroidism, vitamin D deficiency, anemia, iron overload, serological positivity for Hepatitis-C virus and history of blood transfusion are all the factors that have been reported to induce a poor response to HB vaccine in CKD patients.^[7,12,20,34,37,39,41-43]

In our study, age and sex showed an effect in seroconversion rates in univariate analysis. However, in multivariate analysis, there was no significant relationship between sex and seroconversion. In addition, multivariate analysis showed that older patients with the lowest level of kidney function were less likely to seroconvert.

The findings can be summarized as follows. Currently, it is not clear whether the different stages of CKD, as well as age and sex could be an independent predictor of seroconversion after administration of HB vaccine. However, our study found that in earlier stages of kidney disease, vaccination is more likely to induce seroconversion compared with the latest stage, which is the dialysis stage.

Therefore, kidney function appears to be an independent predictor of seroconversion. In addition, older HD patients appeared to be less likely to seroconvert.

We also found that an increased response rate of 89.7% occurred when 40µg doses of vaccine were used on patients with stages 3-4 of CKD, leading us to conclude that more seroconversion may be obtained in such higher vaccine dosages.

Limitation

Of course, a higher number of CKD patients at stages 3-4 could have further strengthened the results of this study.

CONCLUSION

We suggest that as soon as patients are diagnosed with CKD, they should be administered HB vaccine. We also encourage further studies to assess if higher seroprotection rates could be obtained with administration of higher dosage of vaccine at earlier stages of CKD.

ACKNOWLEDGMENTS

The authors would like to thank the kidney dialysis center, nurses, and patients of Vli-e-Asr Hospital in the city of Zanjan.

REFERENCES

1. Janus N, Vacher LV, Karie S, Ledneva E, Deray G. Vaccination and chronic kidney disease. *Nephrol Dial Transplant* 2008;23:800-7.
2. Shepard CW, Simard EP, Finelli L, Fiore AE, Bell BP. Hepatitis B virus infection: Epidemiology and vaccination. *Epidemiol Rev* 2006;28:112-25.
3. CDC. Recommendations of the immunization Practices Advisory Committee (ACIP). Inactivated Hepatitis B virus vaccine. *MMWR Morb Mortal Wkly Rep* 1982;31:317-22.
4. Azarkar Z, Sharifzadeh GH. Efficacy of HBV vaccination in children with thalassemia major, south Khorasan province, Iran. *Iranian Red Crescent Medical Journal* 2009;11:318-20.
5. Merat S, Rezvan H, Nouraei M, Jamali A, Assari S, Abolghasemi H, *et al.* The prevalence of Hepatitis B surface antigen and anti-Hepatitis B core antibody in Iran: A population-based study. *Arch Iran Med* 2009;12:225-31.
6. Poorolajal J, Majdzadeh R. Prevalence of chronic Hepatitis B infection in Iran: A review article. *JRes Med Sci* 2009;14: 249-58.
7. Tang S, Lai KN. Chronic viral hepatitis in hemodialysis patients. *Hemodial Int* 2005;9:169-79.
8. Fabrizi F, Martin P. Hepatitis B virus infection in dialysis patients. *Am J Nephrol* 2000;20:1-11.
9. Finelli L, Miller JT, Tokars JI, Alter MJ, Arduino MJ. National surveillance of dialysis-associated diseases in the United States, 2002. *Semin Dial* 2005;18:52-61.
10. Igaki N, Nakaji M, Moriguchi R, Akiyama H, Tamada F, Oimomi M, *et al.* An outbreak of fulminant hepatitis in immunocompromised hemodialysis patients. *J Gastroenterol* 2003;38:968-76.
11. Tokars JI, Finelli L, Alter MJ, Arduino MJ. National surveillance of dialysis-associated diseases in the United States, 2001. *Semin*

- Dial 2004;17:310-9.
12. Fabrizi F, Bunnapradist S, Martin P. HBV infection in patients with end-stage renal disease. *Semin Liver Dis* 2004;24 (Suppl 1):63-70.
 13. Fehr T, Ambühl PM. Chronic hepatitis virus infections in patients on renal replacement therapy. *Nephrol Dial Transplant* 2004;19:1049-53.
 14. Burdick RA, Bragg-Crescham JL, Woods JD, Hedderwick SA, Kurokawa K, Combe C, *et al.* Patterns of hepatitis B prevalence and seroconversion in hemodialysis units from three continents: The DOPPS. *Kidney Int* 2002;63:22229.
 15. CDC. Recommendations for preventing transmission of infections among chronic haemodialysis patients. *MMWR Recomm Rep* 2001;50:1-43.
 16. Charest AF, McDougall J, Goldstein MB. A randomized comparison of intradermal and intramuscular vaccination against Hepatitis B virus in incident chronic hemodialysis patients. *Am J Kidney Dis* 2000;36:976-82.
 17. Tokars JI, Frank M, Alter MJ, Arduino MJ. National surveillance of hemodialysis associated diseases in the United States, 2000. *Semin Dial* 2002;15:162-71.
 18. Liu YL, Kao MT, Huang CC. A comparison of responsiveness to Hepatitis B vaccination in patients on hemodialysis and peritoneal dialysis. *Vaccine* 2005;23:3957-60.
 19. DaRoza G, Loewen A, Djurdjev O, Love J, Kempston C, Burnett S, *et al.* Stage of chronic kidney disease predicts seroconversion after Hepatitis B immunization: Earlier is better. *Am J Kidney Dis* 2003;42:1184-92.
 20. Leehey DJ, Cannon JP, Lentino JR. *Handbook of dialysis*, John T. Daugirdas, Peter Gerard Blake, Todd S. Ing. 4th ed. 2007 . Lippincott Williams & Wilkins, Dec 8, 2006 - 774 pages.
 21. Yoon JW, Gollapudi S, Pahl MV, Vaziri ND. Naive and central memory T-cell lymphopenia in end-stage renal disease. *Kidney Int* 2006;70:371-6.
 22. Pesanti EL. Immunologic defects and vaccination in patients with chronic renal failure. *Infect Dis Clin North America* 2001;15:813-32.
 23. Department of Health. Good practice guidelines for renal dialysis/transplantation units. Prevention and control of blood-borne virus infection. Recommendation convened by the Public Health Laboratory Service on behalf of the Department of Health. Department of Health Publications; 2002. www.dh.gov.uk
 24. Taheri S, Shahidi S, Moghtaderi J, Seirafian S, Emami A, Eftekhari S. Response rate to Hepatitis B vaccination in patients with chronic renal failure and end-stage-renal- disease: Influence of diabetes mellitus . *Journal of Research in Medical Sciences*, 2005;10:384-90.
 25. Dukes CS, Street AC, Starling JF, Hamilton JD. Hepatitis B vaccination and booster in predialysis patients: A 4-year analysis. *Vaccine* 1993;11:1229-32.
 26. Vlassopoulos D. Recombinant Hepatitis B vaccination in renal failure patients. *Curr Pharm Biotechnol* 2003;4:141-51.
 27. Girndt M, Kohler H. Hepatitis B virus infection in haemodialysis patients. *Semin Nephrol* 2002;22:340-50.
 28. Fabrizi F, Ganeshan SV, Dixit V, Martin P. Meta-analysis: The adjuvant role of granulocyte macrophage-colony stimulating factor on immunological response to Hepatitis B virus vaccine in end-stage renal disease. *Aliment Pharmacol Ther* 2006;24:789-96.
 29. Cruciani M, Mengoli C, Serpelloni G, Mazzi R, Bosco O, Malena M. Granulocyte macrophage colony-stimulating factor as an adjuvant for Hepatitis B vaccination: A meta-analysis. *Vaccine* 2007;25:709-18.
 30. Argani H, Akhtarishojaie E. Levamisole enhances immune responsiveness to intra-dermal and intra-muscular Hepatitis B vaccination in chronic hemodialysis patients. *J Immune Based Ther Vaccines* 2006;4:3.
 31. Levin A. Dialysis: Intradermal HBV vaccination is preferable in non-responders. *Nat Rev Nephrol* 2009;5:616-7.
 32. Barraclough KA, Wiggins KJ, Hawley CM, Van Eps CL, Mudge DW, Johnson DW, *et al.* Intradermal versus intramuscular hepatitis B vaccination in hemodialysis patients: A prospective open-label randomized controlled trial in nonresponders to primary vaccination. *Am J Kidney Dis* 2009;54:95-103. Epub 2009 May 29.
 33. Mat O, Mestrez F, Beauwns R, Muniz-Martinez MC, Dhaene M. Primary high dose intradermal Hepatitis B vaccination in hemodialysis: Cost-effectiveness evaluation at 2 years. *Hemodial Int* 2006;10:49-55.
 34. Kong NC, Beran J, Kee SA, Miguel JL, Sánchez C, Bayas JM, *et al.* A new adjuvant improves the immune response to Hepatitis B vaccine in hemodialysis patients. *Kidney Int* 2008;73:856-62.
 35. Morais EO, Resende MR, Oliveira AM, Sinkoc VM, Garcia MT, Angerami RN, *et al.* Intradermal Hepatitis B vaccination in patients with advanced chronic renal failure: Immunogenicity and follow-up. *Aliment Pharmacol Ther*. 2007;25:849-55.
 36. Fabrizi F, Ganeshan SV, Dixit V, Martin P. Meta-analysis: Intradermal vs. intramuscular vaccination against Hepatitis B virus in patients with chronic kidney disease. *Aliment Pharmacol Ther* 2006;24:497-506.
 37. McNulty CA, Bowen JK, Williams AJ. Hepatitis B vaccination in predialysis chronic renal failure patients a comparison of two vaccination schedules. *Vaccine* 2005;23:4142-7.
 38. Siddiqui S, Malik A, Shukla I, Rizvi M, Haque SF. Seroprotection after Hepatitis B vaccination in chronic kidney disease patients with modified schedule and dosage. *J Infect Dev Ctries* 2010; 4:389-92.
 39. Fraser GM, Ochana N, Fenyves D, Neumann L, Chazan R, Niv Y, *et al.* Increasing serum creatinine and age reduce the response to hepatitis B vaccine in renal failure patients. *J Hepatol* 1994;21:450-4.
 40. Watkins SL, Alexander SR, Brewer ED, Hesley TM, West DJ, Chan IS, *et al.* Response to recombinant Hepatitis B vaccine in children and adolescents with chronic renal failure. *Am J Kidney* 2002;40:365-72.
 41. Kara IH, Yilmaz ME, Suner A, Kadiroglu AK, Isikoglu B. The evaluation of immune responses that occur after HBV infection and HBV vaccination in hemodialysis patients. *Vaccine* 2004;22:3963-7.
 42. Fabrizi F, Martin P, Dixit V, Bunnapradist S, Dulai G. Meta-analysis: The effect of age on immunological response to Hepatitis B vaccine in end-stage renal disease. *Aliment Pharmacol Ther* 2004;20:1053-62.
 43. Fisman DN, Agrawal D, Leder K. The effect of age on immunologic response to recombinant Hepatitis B vaccine: A meta-analysis. *Clin Infect Dis* 2002;35:1368-75.

How to cite this article: Ghadiani MH, Besharati S, Mousavinasab N, Jalalzadeh M. Response rates to HB vaccine in CKD 34 and hemodialysis patients. *J Res Med Sci* 2012;17:527-33

Source of Support: Nil, **Conflict of Interest:** None declared.